HARRISON COUNTY: Key Data Points

- Compared to the state, a higher percentage of Harrison county mothers are utilizing prenatal care during the first trimester of pregnancy (88.8% versus 80.6% respectively), but a higher percentage are also delivering preterm (10.3% versus 9.9% respectively)
- Women in the county are breastfeeding less on discharge (51.4% versus the state's 63.2%)
- Harrison county observed 21 accidental deaths (12 of which were motor vehicle), and 1 suicidal death

Note: All figures relate to 2004 data from the Indiana State Department of Health, unless otherwise specified

Figure 1: Births

	Total		White		Black		Other	
	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*
Harrison County	435	12.0	430	12.0	2	11.8(U)	3	8.1(U)
INDIANA	87,125	14.0	75,680	13.7	9,582	17.5	1,863	11.7

^{*} births per 1,000 population; U = unstable rate, based on fewer than 20 births

Figure 2: Infant Mortality

			NUI	MBER			RATE*	
COUNTY OF	REPORTED RACE/	Ling	Infant Deaths (< 1 yr)	Neonatal Deaths (< 28 days)	Post Neonatal Deaths	Infant Deaths (< 1 yr)	Neonatal Deaths (< 28 days)	Post Neonatal Deaths
RESIDENCE	ETHNICITY	Live Births	(4.7.)	(120 am jo)	(28-364 days)	(- , , ,	(20 aa) 0,	(28-364 days)
HARRISON	All Races	435	3	3	0	**	**	0.0
	White	430	3	3	0	**	**	0.0
	Black	2	0	0	0	0.0	0.0	0.0
	Other	3	0	0	0	0.0	0.0	0.0
	Hispanic	8	0	0	0	0.0	0.0	0.0
INDIANA	All Races	87,125	702	472	230	8.1	5.4	2.6
	White	75,680	522	354	168	6.9	4.7	2.2
	Black	9,582	164	105	59	17.1	11.0	6.2
	Other	1,863	16	13	3	**	**	**
	Hispanic	7,237	65	50	15	9.0	6.9	**

^{*}rate per 1,000 live births **Numerator is less than 20 and the rate is unstable

Figure 3: Outcome Indicators of Mother, by Race


Birth '		/eight				During Pregnancy		
HARRISON	Total	% Low	% Very Low (< 1500	% Preterm	% PNC 1st	%	%	% Unmarried
COUNTY	Births	(< 2500 grams)		t	Trimester ‡	Alcohol	Smoked	Parents
White	430	6.3	0.2	9.1	73.7(S)	0.2	24.4	33.3
Black	2	**	**	**	**	**	**	**
All Races	435	6.2	0.2	9.2	73.6	0.2	24.4	33.1


		Birth Weight				During Pregnancy		
	Total	% Low	% Very Low	% Preterm	% PNC 1st	%	%	% Unmarried
INDIANA	Births	(< 2500 grams)	(< 1500 grams)	†	Trimester [‡]	Alcohol	Smoked	Parents
All Races	87,125	8.1	1.5	10.2	79.3	0.6	18.0	38.8
White	75,680	7.4	1.3	9.9	80.8	0.6	18.8	34.4
Black	9,582	13.7	3.0	13.9	67.1	0.9	14.4	77.9


[†] Preterm: less than 37 weeks gestation

Figure 4: Breastfeeding

HARRISON COUNTY	All Races	White	Black
Total Births	435	430	2
Breastfeeding on Discharge	216	213	2
% Breastfeeding on Discharge	49.7	49.5	100.0







[‡] PNC: Prenatal Care 'S' Significantly different than the state percent

^{&#}x27;**' Outcomes for this county are suppressed due to 20 or fewer total births

Figure 5: HIV/AIDS

	Total # of AIDS cases reported since 1982	Total # of HIV cases reported since 1985
HARRISON	11	8
COUNTY	7,000	2.045
INDIANA	7,989	3,945

^{*}numbers current through June 2006

Figure 6: Nutritional Risk Factors by Status,

Indiana WIC Clinics, Quarter Ending 09/30/2004

Local Agency: Blue River Services, Inc Counties Included: Harrison, Washington								
	Childr	en 1-5 y			%			
Risk Factor or Status	White	Black	Hispanic	Asian	Other	Total	within Agency	
Overweight	38	0	2	1	0	41	5.4%	
At Risk for becoming Overweight	39	0	1	0	0	40	5.2%	
	Local Agency: State Wide Counties Included: All							
	Childr	en 1-5 y	ears of Ago	e			%	
Risk Factor or Status	White	Black	Hispanic	Asian	Other	Total	within Agency	
Overweight	1,732	445	800	12	97	3,086	4.8%	
At Risk for								

639

485

2,023

becoming Overweight 3,282

5.1%

124

11

Priority Benchmarks (Harrison & Inc	diana)	_
Demographics: Percentage of population enrolled in Medicaid, Percentage of population enrolled in Food Stamps.	Harrison	Indiana
1 % of Population enrolled in Medicaid as of June 2003	10.7	12.4
2 % of Population in Food Stamps Program, 2005	7.0	9.0
Priority #1 (pregnancy outcomes): Average Infant Mortality Rate, Average Fetal death Rate	Harrison	Indiana
1 Av Total Infant Death Rate (1995-2004)	5.6	7.9
2 Average Fetal Death Rate (1995-2004)	8.4	6.6
Priority #2 (reduce barriers): Health Professional Shortage Areas, Percentage of women with prenatal care in first trimester	Harrison	Indiana
1 Scores for Primary Care Health Professional Shortage Areas (HPSAs) (2005)	8	NA
2 % PNC 1st Trimester†, 2004	73.6	79.3
Priority #4 (environmental): Prevalence of asthma, Childhood lead poisoning	Harrison	Indiana
Prevalence of Asthma in a Medicaid Cohort of Children age 0-17 years, 2003	9.2	10.4
2 %Estimated Lead Blood Levels (%EBLL) for Children under 6 years old FY 2005-2006	2.42	2.42
Priority #5 (tobacco): Percentage of mothers who smoked during pregnancy, Percentage of Adult Smoking	Harrison	Indiana
1 % of mothers Smoked During Pregnancy, 2004	24.4	18.0
2 % of Adult Smoking in 2003	31.4	29.3
Priority #7 (adolescent risk behaviors): Adolescent deaths due to motor vehicle accidents, Chlamydia rates (ages 15-24).	Harrison	Indiana
Number of Adolescent (15-24) deaths due to Motor vehicle accidents (2002-2004)	11	771
2 Chlamydia rates per 100,000 county specific population (15-24yrs of Age)	614.75	1602.35
Priority #8 (obesity): Obesity in pregnant women	Harrison	Indiana
Number of Obese Pregnant Women In Indiana, 2004	19	2,827
Priority #9 (domestic violence, childhood injury): Child deaths due to Injury, Child abuse and neglect rates	Harrison	Indiana
1 Child deaths due to Injury 1-14yrs (2002-2004)	1	279
2 Child Abuse and Neglect per 1,000 Population under age 18, 2004	18.2	13.5
Priority #10 (To improve racial and ethnic disparities in women of childbearing age, mothers, and children's health outcomes): Percentage of Hispanic Population of any Race, LBW Disparity Ratio	Harrison	Indiana
1 % of Hispanic Population of any Race, 2005	1.4	4.5
2 Low Birth Weight Disparity Ratio (B/W) (1995-2004)	6.89	1.88

Source: Data Analysis Team, ISDH; Health Resources and Service Administration; Indiana Prevention Resource Center; Data analysis team, Office of Medicaid Policy & Planning. Notes: 1. NS= Not Stable 2. NA = Not Available 3. Infant Mortality rate and Fetal Death rate is for 1000 live births. 4. Low Birth Weight: Birth Weight Less than 2,500grams. 5. HPSA Score ranges 1 – 25. The higher the score, the higher the need. 6. EBLL: Ten (10) micrograms of lead per deciliter of whole blood for one venous test; %EBLL is based on screening rate of 27% Statewide, unduplicated kids 7. Obesity in Pregnant Women determined by "Concurrent Illness Obesity" checkbox on the birth certificate 8. PNC percentages are calculated using total births in each County.